

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 756 828 B1

(12)

EUROPEAN PATENT SPECIFICATION(45) Date of publication and mention
of the grant of the patent:
18.11.1998 Bulletin 1998/47(51) Int. Cl.⁶: **A23L 1/308**, A23L 1/09,
A23L 1/05, A61K 31/715(21) Application number: **95202142.6**(22) Date of filing: **04.08.1995****(54) Nutritional composition containing fibres**

Diätfasern enthaltende Nahrungszusammensetzung

Composition nutritionnelle contenant des fibres diététiques

(84) Designated Contracting States:
BE CH DE ES FR GB GR LI NL SE(43) Date of publication of application:
05.02.1997 Bulletin 1997/06(73) Proprietor: N.V. Nutricia
NL-2712 HM Zoetermeer (NL)(72) Inventors:
• Green, Celrwen Jane
NL-2716 NN Zoetermeer (NL)
• Bork, Roelof André
NL-2727 GR Zoetermeer (NL)
• Hageman, Robert Johan Joseph
NL-2742 EV Waddinxveen (NL)• Boerma, Jan Albertus
NL-2723 AA Zoetermeer (NL)
• Van Hoey-De Boer, Klaske Anna
NL-3443 BD Woerden (NL)(74) Representative:
de Bruijn, Leendert C. et al
Nederlandsch Octrooibureau
P.O. Box 29720
2502 LS Den Haag (NL)(56) References cited:
EP-A- 0 483 070 EP-A- 0 486 425
EP-A- 0 504 055 EP-A- 0 570 791
WO-A-90/04334 WO-A-93/00020
DE-A- 2 937 358 US-A- 5 229 136

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 0 756 828 B1

Description

The invention relates to a dietary fibre composition which can be used in various types of nutritional compositions, such as clinical food, infant food or dietary food.

Fibre plays an important role in the nutrition of healthy people. It maintains gut function and clears toxic compounds, by providing stool bulk and substrate to intestinal flora, and keeps the gut wall in good condition. Hospital patients who are critically ill or chronically sick or comatose patients who normally receive enteral clinical nutrition (ECN), need fibre for these purposes, and consumption of the right fibre mixture is especially important for patients who suffer from intestinal problems, such as ulcerative colitis, Crohn's disease, and the like, and who have received antibiotics. Also persons with constipation or with diarrhoea have special interest in this kind of nutrition, and the same applies to persons who are not able to consume the recommended daily amount of fibre for whatever reason (e.g. diet).

An optimal fibre composition will:

- be similar to the normal fibre composition as consumed by healthy persons in Western countries;
- produce beneficial short-chain fatty acids (SCFA) in a balanced ratio and at a sufficient and uniform rate during transit through the ileum and the large intestine;
- not produce gas at too high or too low levels;
- give sufficient mass to the faeces to facilitate defecation;
- not drastically increase the viscosity of the faeces.

A dietary fibre composition for use in nutritional products is disclosed in International patent application WO 93/00020. It contains 5-50 wt.% of soluble, fermentable fibre (gum arabic), 5-20 wt.% of soluble, non-fermentable fibre (carboxymethyl-cellulose), and 45-80 wt.% of insoluble, non-fermentable fibre (oat hull). A drawback of this prior art fibre composition is that it does not show an optimal performance as to uniformity and type of short-chain fatty acids and gas production. Also, it does not result in sufficient stool bulk for hospitalised patients. Furthermore, it contains a non-natural, chemically modified fibre, which is not desired in most nutritional compositions. It does not resemble average fibre consumption patterns in Western countries.

EP-A-483070 discloses nutritional compositions containing 10-60 g per day of hydrolysed soluble fibre (hydrolysed pectin and hydrolysed guar). EP-A-486425 discloses food compositions having a pH of 3.5-3.9 containing 0-17 energy % of fibre. EP-A-504055 describes dietetic and therapeutic compositions containing modified starch and soluble fibres (pectin or soy fibres). EP-A-570791 discloses enteral nutritional low-calorie compositions for tube-fed patients with diarrhoea, containing 20 g/l of fibre (carob extract, pectin and soy polysaccha-

ride). These products do not provide a balanced fibre composition as referred to above.

The object of the present invention is to provide a dietary fibre mixture which meets the requirements of an optimal composition as indicated above, and to provide nutritional compositions suitable for enteral administration, containing such a fibre mixture.

The composition according to the invention is characterised in that it contains 5-120 g of fibre per daily dosage of the composition, and the fibre consists of 15-50 wt.% of soluble non-starch polysaccharides, 15-45 wt.% of insoluble non-starch polysaccharides, and 8-70 wt.% of oligosaccharides and/or resistant starch.

The novel fibre mixture is fermented at a relatively uniform rate, starting in the first part of the colon and continuing in more distal parts of the colon, to yield a relatively constant amount of SCFA's and a relatively high amount of propionic and/or butyric acid and to produce a relatively constant level of gas. Part of the mixture will not be fermented and passes to subsequent parts of the intestine, where it will contribute to the stool bulk and will also not increase substantially the viscosity of the faeces.

In the context of this specification, enteral administration includes liquid and dry administration forms. Liquid compositions for enteral administration comprise tube-feeds (via nasogastric, gastrostomy, jejunostomy tubes), orally consumed compositions (drinks), both as complete foods and as supplements. Dry compositions comprise powders, e.g. in cans or sachets, which are to be reconstituted or to be added to other food compositions, and also bars and the like.

The term "fibre" or "dietary fibre" is used herein to denote plant-derived food material, in particular oligo- and polysaccharides (cellulose and hemicellulose), lignin, and resistant starch, that is not digested by the human (non-bacterial) enzymes of the intestinal tract. This means that fibre is that part of the food that is not absorbed in the small intestine and thus enters the large intestine (colon).

The term "fermentable fibre" means fibre that undergoes (anaerobic) breakdown by microorganisms in the large intestine to smaller molecules, in particular to short-chain fatty acids, such as acetic, propionic and butyric acid. Fermentation also results in the production of gases, such as carbon dioxide, hydrogen and methane. Fibre fermentability may be determined e.g. by the method described in Amer. J. Clinical Nutrition (1991), 53, 1418-1424.

The term "soluble fibre" means that the fibre is at least 50% soluble according to the method described by L. Prosky et al., J. Assoc. Off. Anal. Chem. 71, 1017-1023 (1988).

Examples of soluble non-starch polysaccharides include inulin (a polysaccharide mainly consisting of fructose units), pectin (a polysaccharide mainly consisting of galacturonic acid and rhamnose), β -glucans (small cellulose-type branched glucose polymers), gum

arabic, tragacanth, mucilages, guar and locust bean gum, agar, carageenans, alginates, xanthan and the like. Most of these soluble fibres are fermentable for the largest part. In particular the proportion of soluble non-starch polysaccharides is within the range of 20-40 wt.%. 5

Examples of insoluble non-starch polysaccharides include cellulose (for example derived from oat hull, soybeans, cereal bran) and hemicellulose (mostly branched arabinoxylans or galactans, e.g. from cereals, potatoes or soybeans). Most of these insoluble fibres are partly fermentable or non-fermentable. In particular the proportion of insoluble non-starch polysaccharides is within the range of 20-40 wt.%. 10

In addition to the non-starch polysaccharides (soluble and insoluble), the fibre composition according to the invention contains either oligosaccharides or resistant starch or preferably both. It may also contain lignin. Oligosaccharides are understood to comprise any saccharide containing at least two and up to 20 monosaccharide units, whether of starch (α -glucan) or non-starch type. Examples of non-digestible oligosaccharides include fructo- and galacto-oligosaccharides, such as hydrolysed inulin and levan (fructans), and short-chain amylopectins and the like. The oligosaccharides are usually soluble and largely fermentable. Especially preferred are oligosaccharides having an average chain length (degree of polymerisation) of between 3 and 9 monosaccharide units. Preferably, the fibre composition according to the invention consists of 8-40 wt.%, more preferably of 12-30 wt.% of oligosaccharides, especially hydrolysed inulin, or a derivative thereof. 20

Resistant starch is commonly known and generally consists of any type of starch that escapes digestion and intestinal absorption and can be formed during processing of starchy foods. In some starches the crystalline structure will be altered or retrograded by these processes, which make the starch difficult to be digested. Resistant starch may also be obtained from raw foods, such as potatoes or green bananas. Resistant starch is normally insoluble and the fermentability varies with the source. Preferably, resistant starch is present in the fibre mixture of the invention in addition to the oligosaccharides, especially at a level of 5-30 wt.%, more preferably of 8-25 wt.% with respect to the total fibre composition. 30

When all components of the fibre mixture according to the invention are classified as soluble or insoluble, the fibre consists of 25-80 wt.% of soluble fibre and 20-75 wt.% of insoluble fibre. The soluble part of the fibre mixture, mainly comprising oligosaccharides and soluble non-starch polysaccharides, and some of the resistant starch, should have a fermentability of at least 80%, preferably at least 90%. In particular, the fibre mixture consists of 35-65 wt.% of soluble fibre. The insoluble part of the fibre mixture, mainly comprising resistant starch and insoluble non-starch polysaccharides (cellulose, hemicellulose), should have a fermentability of 35

between 5 and 50%, especially between 20 and 40%. In particular, the fibre mixture contains 35-65 wt.% of insoluble fibre.

In addition to the fibre polysaccharides, the nutritional composition according to the invention also preferably contains small amounts of lignin or lignin derivatives, in particular, 1-5 wt.% of lignin. Preferably, the composition does not contain chemically modified fibres, such as carboxymethyl starch of hydroxyethyl cellulose.

The fibre content of the nutritional compositions according to the invention can be expressed in the amount of fibre to be taken per day, as defined above, i.e. 5-120 g per daily dosage, or especially 8-70 g per daily dosage. Instead, the fibre content for non-low-calorie compositions can also be expressed as weight of fibre per energy unit. Thus, the nutritional composition may contain 5-120 g, preferably 8-70 g of fibre per 2000 kcal of energy. These values apply both to dry compositions and to liquid compositions or reconstituted compositions. 40

For dry compositions, it may be convenient to express the fibre content with respect to the total dry matter of the composition. Thus dry nutritional compositions according to the invention may contain 0.5-100 g of fibre per 100 g. The higher levels, e.g. 50-100 g per 100 g, apply to dry fibre supplements to be administered in combination with other, standard food compositions. An example is a 100% fibre powder. Intermediate levels, e.g. between 5 and 80 g per 100 g, may apply to fibre-reinforced food supplements or complete nutritional compositions, e.g. for clinical uses. As an example, a low-calorie bar may contain 20-80 g of fibre per 100 g of product, the remainder consisting of e.g. maltodextrin and/or protein. The lower levels, e.g. between 0.5 and 25 g per 100 g, may apply to special compositions, such as infant formulae. For example, a dry infant formula can contain 1 g of fibre in a total of 100 g. 50

As an alternative, the nutritional composition may be a liquid composition, or a powder which is to be reconstituted by dilution with water. In that case, the composition contains 5-120 g of fibre per l, in particular 10-100 g/l. In case of liquid compositions, the pH is preferably between 4 and 8.

Apart from a pure fibre supplement, the nutritional composition will further contain digestible carbohydrates and/or fats and/or proteinaceous material.

The digestible carbohydrates (to be distinguished from the indigestible fibre carbohydrates) constitute the energy source in most cases. Digestible carbohydrates to be used may include monosaccharides such as glucose, fructose, galactose and ribose, disaccharides such as lactose, sucrose and maltose, and digestible oligo- and polysaccharides such as starch-derived material, including maltodextrins. The latter are the preferred carbohydrates for nutritional compositions intended for clinical nutrition.

The fats optionally contained in the nutritional com-

positions may be the usual nutritional fats. Fat-containing compositions according to the invention further preferably contain fats with long-chain polyunsaturated fatty acids (PUFA's), especially parent PUFA's linoleic acid (ω -6 series) and α -linolenic acid (ω -3 series) and higher homologues produced by alternate desaturation and elongation of the parent PUFA's. These higher homologues include γ -linolenic acid (GLA: C18:3 ω -6) and stearidonic acid (SDA: C18:4 ω -3) and subsequent homologues (C20:3, C20:4, C22:4, C22:5, etc. of the ω -6 series, and C20:4, C20:5, C22:5, etc. of the ω -3 series). Including the parent PUFA's, the preferred level is between 3.5 and 12 energy% of the composition, especially 5-9 en.%. The preferred level of the total homologues (GLA + SDA and higher) is between 0.2 and 5 en.%, especially between 0.5 and 2.5 en.% of the composition. Preferably, a fat-containing composition also contains an antioxidant such as tocopherol.

The proteins to be optionally used in the nutritional compositions according to the invention can be those commonly present in tube feeding, infant formulae and other foodstuffs. Preferred proteinaceous materials include caseins, whey, soy protein, and partially denatured whey proteins, or partially hydrolysed whey proteins. The partly denatured whey proteins are especially beneficial for complete nutritional compositions. The preferred level of partly denatured whey proteins corresponds to at least 20 % of the total protein content, more preferably 40-100 wt.% of the total protein content, or alternatively preferably 2-30, more preferably 4-20 wt.% of the dry matter content of the composition.

Example 1

A fibre composition was prepared by mixing 32 wt.% of soy polysaccharide, 12 wt.% of α -cellulose, 24 wt.% of gum arabic, 23 wt.% of Ratiolose® (hydrolysed inulin) and 9 wt.% of Novelose® (resistant starch), followed by spray-drying.

Example 2

A complete liquid clinical food composition was prepared by mixing 1-2 g of the fibre mixture according to example 1, 4-25 energy% (1.0-7.5 g) of protein, 40-78 en.% (12-24 g) of carbohydrate, 8-50 en.% (1-11 g) of fat including 0.5-1.5 g of PUFA's, in addition to Na (90 mg), K (180 mg), Cl (130 mg), Ca (85 mg), P (70 mg) and Mg (26 mg) per 100 g, and further Fe (1.2 mg), Zn (1.2 mg), Cu (0.2 mg), Mn (0.3 mg), F 0.1 μ g, Mo (6 μ g), Se (6 μ g), Cr (4 μ g), I (10 μ g), vitamins (A, D, E, K, B6, B12, C, thiamine, riboflavin, niacin, pantothenic acid, folic acid, biotin), choline (15-70 mg) and myo-inositol (15-40 mg) per 100 ml of water.

Claims

1. Nutritional composition, which, optionally after

admixture of other food components and/or dilution with water, is suitable for enteral administration, comprising dietary fibre, characterised in that the fibre consists of 15-50 wt.% of soluble non-starch polysaccharides, 15-45 wt.% of insoluble non-starch polysaccharides, and 8-70 wt.% of non-digestible oligosaccharides and/or resistant starch.

2. Nutritional composition according to claim 1, wherein the fibre comprises 20-40 wt.% of soluble non-starch polysaccharides and 20-40 wt.% of insoluble non-starch polysaccharides.
3. Nutritional composition according to claim 1 or 2, wherein the fibre comprises 8-40 wt.%, preferably 12-30 wt.% of non-digestible oligosaccharides.
4. Nutritional composition according to any one of claims 1-3, wherein the non-digestible oligosaccharides comprise hydrolysed inulin or a derivative thereof.
5. Nutritional composition according to any one of claims 1-4, wherein the non-digestible oligosaccharides have all average degree of polymerisation of 2 to 20 mono-saccharide units.
6. Nutritional composition according to any one of claims 1-5, wherein the fibre comprises 5-30 wt.%, preferably 8-25 wt.% of resistant starch.
7. Nutritional composition according to any one of claims 1-6, which contains 5-120 g of fibre per 2000 kcal of energy.
8. Nutritional composition according to any one of claims 1-7, which is a dry composition and contains 0.5-100 g of fibre per 100 g.
9. Nutritional composition according to any one of claims 1-8, which is a liquid composition, optionally after reconstitution, and contains 5-120 g of fibre per l.
10. Nutritional composition according to any one of claims 1-9, wherein the composition does not contain chemically modified fibres.
11. Nutritional composition according to any one of claims 1-10, wherein the composition further contains digestible carbohydrates and/or fats and/or proteinaceous material.
12. Nutritional composition according to any one of claims 1-11, further comprising 1-5 wt.% of lignin.

Patentansprüche

1. Nahrungsmittelzusammensetzung, die gegebenenfalls nach Vermischen mit anderen Nahrungsmittelkomponenten und/oder nach Verdünnung mit Wasser sich zur enteralen Verabreichung eignet, enthaltend diätetische Fasern, dadurch gekennzeichnet, daß die Fasern aus 15-50 Gew.-% löslichen Nichtstärke-Polysacchariden, 15-45 Gew.-% unlöslichen Nichtstärke-Polysacchariden und 8-70 Gew.-% nicht-verdaubaren Oligosacchariden und/oder resistenter Stärke bestehen.
2. Nahrungsmittelzusammensetzung nach Anspruch 1, wobei die Fasern 20-40 Gew.-% lösliche Nichtstärke-Polysaccharide und 20-40 Gew.-% unlösliche Nichtstärke-Polysaccharide umfassen.
3. Nahrungsmittelzusammensetzung nach Anspruch 1 oder 2, wobei die Fasern 8-40 Gew.-% und vorzugsweise 12-30 Gew.-% nicht-verdaubare Oligosaccharide umfassen.
4. Nahrungsmittelzusammensetzung nach einem der Ansprüche 1-3, wobei die nicht-verdaubaren Oligosaccharide hydrolysiertes Inulin oder ein Derivat davon umfassen.
5. Nahrungsmittelzusammensetzung nach einem der Ansprüche 1-4, wobei die nicht-verdaubaren Oligosaccharide einen durchschnittlichen Polymerisationsgrad von 2 bis 20 Monosaccharideinheiten aufweisen.
6. Nahrungsmittelzusammensetzung nach einem der Ansprüche 1-5, wobei die Fasern 5-30 Gew.-% und vorzugsweise 8-25 Gew.-% resistente Stärke umfassen.
7. Nahrungsmittelzusammensetzung nach einem der Ansprüche 1 bis 6, enthaltend 5-120 g Fasern pro 2000 kcal Energie.
8. Nahrungsmittelzusammensetzung nach einem der Ansprüche 1 bis 7, bei der es sich um eine trockene Zusammensetzung handelt, die 0,5-100 g Fasern pro 100 g enthält.
9. Nahrungsmittelzusammensetzung nach einem der Ansprüche 1 bis 8, wobei es sich um eine gegebenenfalls nach Rekonstitution erhaltene flüssige Zusammensetzung handelt, die 5-120 g Fasern pro Liter enthält.
10. Nahrungsmittelzusammensetzung nach einem der Ansprüche 1 bis 9, wobei die Zusammensetzung keine chemisch modifizierten Fasern enthält.

11. Nahrungsmittelzusammensetzung nach einem der Ansprüche 1 bis 10, wobei die Zusammensetzung ferner verdaubare Kohlenhydrate und/oder Fette und/oder proteinhaltige Materialien enthält.

12. Nahrungsmittelzusammensetzung nach einem der Ansprüche 1 bis 11, ferner umfassend 1-5 Gew.-% Lignin.

Revendications

1. Composition nutritionnelle, qui, éventuellement après mélange avec d'autres composants alimentaires et/ou dilution avec de l'eau, convient à une administration entérale, comprenant des fibres diététiques, caractérisée en ce que les fibres sont constituées de 15-50 % en poids de polysaccharides solubles autres que l'amidon, 15-45 % en poids de polysaccharides insolubles autres que l'amidon, et 8-70 % en poids d'oligosaccharides non-digestibles et/ou d'amidon résistant.
2. Composition nutritionnelle selon la revendication 1, dans laquelle les fibres comprennent 20-40 % en poids de polysaccharides solubles autres que l'amidon et 20-40 % en poids de polysaccharides insolubles autres que l'amidon.
3. Composition nutritionnelle selon la revendication 1 ou 2, dans laquelle les fibres comprennent 8-40 % en poids, de préférence 12-30 % en poids d'oligosaccharides non digestibles.
4. Composition nutritionnelle selon l'une quelconque des revendications 1-3, dans laquelle les oligosaccharides non digestibles comprennent de l'inuline non hydrolysée ou l'un de ses dérivés.
5. Composition nutritionnelle selon l'une quelconque des revendications 1-4, dans laquelle les oligosaccharides non digestibles ont un degré moyen de polymérisation de 2 à 20 unités monosaccharides.
6. Composition nutritionnelle selon l'une quelconque des revendications 1-5, dans laquelle les fibres comprennent 5-30 % en poids, de préférence 8-25 % en poids d'amidon résistant.
7. Composition nutritionnelle selon l'une quelconque des revendications 1-6, qui contient 5-120 g de fibres pour 2000 kcal d'énergie.
8. Composition nutritionnelle selon l'une quelconque des revendications 1-7, qui est une composition sèche et contient 0,5-100 g de fibres pour 100 g.
9. Composition nutritionnelle selon l'une quelconque des revendications 1-8, qui est une composition

liquide, éventuellement après reconstitution, et contient 5-120 g de fibres par l.

10. Composition nutritionnelle selon l'une quelconque
des revendications 1-9, dans laquelle la composition
ne contient pas de fibres chimiquement modifiées. 5
11. Composition nutritionnelle selon l'une quelconque
des revendications 1-10, dans laquelle la composition
comprend en outre des glucides et/ou des
matières grasses et/ou des substances protéiques
digestibles. 10
12. Composition nutritionnelle selon l'une quelconque
des revendications 1-11, comprenant en outre 1-5
% en poids de lignine. 15

20

25

30

35

40

45

50

55